



Advice Following Implant Placement

Cornwall's Advanced Dental Clinic providing complex treatments like dental implants and cosmetic dentistry every day by highly skilled clinicians backed up by a fantastic qualified dental team.

For more information
Visit puredentalhealth.co.uk
Call 01872 222404



PURE
- dental health -

Advice Following Implant Placement

You will always be given verbal and written **aftercare instructions** when you have dental implant surgery at **Pure Dental Health**, however, should you forget what you've been told or lose your written instructions you can check that you're doing all that you can to ensure a speedy and comfortable recovery by following the advice below.

- Take care to only have cold drinks and do not eat until the local anaesthetic has worn off. Avoid hot drinks or hot food for the first couple of days and do not 'swill' liquid over the area. Try not to disturb the area with your tongue or fingers.
- Soft food and a high protein diet for the first week after surgery will help with the healing process..
- Do not undertake strenuous exercise for the first week after surgery (running / gym).
- You may have some swelling and/or bruising following your treatment - this usually reaches a peak 2 to 3 days later. This is quite normal and both will subside naturally after a few days. Swelling can be reduced with ice packs (or a bag of frozen peas wrapped in a towel). Hold on the cheek area for a maximum of 10 minutes at a time with 20 minutes break. After 24 hours gentle heat is more beneficial. Sleeping propped up slightly on 2-3 pillows may also help.

Advice Following Implant Placement

- Pain should not be a big problem. Although you may be sore for a few days after any surgery in your mouth, this can be easily managed with simple painkillers. You should take these regularly at the maximum stated dose for the first 2-3 days after your surgery. Take whatever painkillers you normally take for headaches, aches and sprains (ibuprofen and paracetamol make a good combination), and take your first dose before the local anaesthetic has completely worn off..
- If after a few days you experience increasing pain and swelling, you must return to [Pure Dental Health](#) as soon as possible so that we can ensure you are not beginning to develop an infection.
- If you have a [denture](#) that covers the surgical area please wear it as little as possible for the first week to protect the surgical site during its initial healing period. You should always leave the denture out at night.
- The stitches are dissolvable but often remain for around 2-3 weeks, if they are uncomfortable or annoying, you may contact us to remove them.
- Some minor bleeding after surgery in the mouth is normal. If this persists, apply pressure by biting down firmly over the area on a dampened gauze swab or clean handkerchief for 60 minutes whilst sitting upright. Do not keep checking or changing the gauze. You should contact us if bleeding persists for any reason after applying pressure in this way.
- If you have been given a course of antibiotics to take after your surgery, please ensure that you complete the course.

Advice Following Implant Placement

Cleaning Advice

- Successful **oral surgery** depends on keeping the mouth as clean as possible.
- Please start to use the mouthwash that you have been given the following morning of your surgery and continue to use for 1 week, twice a day – the same time that you **brush your teeth**. This is very important. (The mouthwash can be purchased from the practice and posted to you for convenience).
- You should also start cleaning your other teeth as normal with a toothbrush, starting on the evening of your surgery. Avoid the surgical site until you see the dentist for your suture removal appointment which will be roughly a week after surgery.
- Hot salt mouthwashes (a cup of hot water with a teaspoon of salt) are very beneficial for healing in the first week but taste awful. Ensure that the mouthwash is not so hot that it scalds. Hold the mouthwash over the surgical site until it cools. Repeat as often as possible.
- Try to keep food away from the surgical area for as long as possible. Gently rinse your mouth after eating to keep the area clean.
- You are advised not to smoke until the wound has healed as this severely limits healing in the mouth.

Advice Following Implant Placement

- We want your recovery to be as smooth and pleasant as possible. It is vital to follow these instructions very carefully - if you have any concerns or questions regarding your progress, please do not hesitate to **contact us**. An out of hours contact number is always available on our answer phone.